

[GNFAC Avalanche Forecast for Mon Jan 2, 2023](#)

Good morning. This is Alex Marienthal with the Gallatin National Forest Avalanche Forecast on Monday, January 2nd at 6:50 a.m. This information is sponsored by [Spark R&D](#) and [onX](#). This forecast does not apply to operating ski areas.

AVALANCHE FATALITY

On December 31, 2022 a snowmobiler was killed in a very large avalanche north of Cooke City. One rider in a group of two triggered the avalanche, was carried 600 vertical feet, and buried 5 feet deep. He was located with a [probe](#) line an hour after the avalanche happened and was unable to be revived. Our deepest condolences go out to the family, friends, and those involved with the rescue and recovery ([Full details and photos here](#)).

Mountain Weather

This morning there is no new snow, temperatures are single digits to teens F, and wind is out of the east to northeast at 0-10 mph. In the Bridger Range wind is east at 20 mph with gusts to 35 mph. Today will be mostly sunny with temperatures in the teens to low 20s F and light to moderate easterly wind.

Snowpack and Avalanche Discussion



[Bridger Range](#) [Gallatin Range](#) [Madison Range](#) [Lionhead Range](#) [Cooke City](#)

A person can trigger large to very large avalanches breaking on buried weak layers. The avalanche that killed a snowmobiler on Saturday near Cooke City broke 1.5-4 feet deep on a layer of sugary, faceted snow below a very hard slab ([video of site investigation](#)). Snow from the past week was drifted onto the slope to further stress the weak layer towards breaking. Also, a snowmobiler triggered slide on Sheep Mountain near Cooke City broke 1-4 feet deep ([video](#), [details](#)), and another on Lionhead ridge near West Yellowstone broke 2-4 feet deep ([details](#)). All these avalanches broke on buried persistent weak layers which means similar avalanches continue to be possible until the snowpack has an extended break from the stress of new and wind-drifted snow.

Avalanches of recently wind-drifted snow are also possible to trigger. These could be large enough to bury a person, and especially dangerous if they push you into hazards like trees, rocks, or over cliffs. Yesterday, skiers north of Bridger Bowl changed their plans after finding unstable snowpack test results below a slab of wind-drifted snow ([observation](#)), and there was a natural avalanche on Saddle Peak that involved recent wind-drifted snow ([photo](#)). Today's east winds will form drifts in less common locations.

The snowpack depth is shallower in northern areas than further south, but the list of avalanche concerns and travel advice is the same. Carefully assess the snowpack on each steep slope you plan to ride or cross. Watch for signs of unstable drifts such as cracking across the snow surface and avoid steep slopes where you suspect buried weak layers exist. Slopes with a relatively shallower snowpack or where the snow depth varies greatly across a slope are the most suspect and first to be avoided, but you should dig and probe to look for buried weak layers before riding or crossing any steep slope. Today, buried weak layers and recently drifted snow make human-triggered avalanches possible and the avalanche danger is [MODERATE](#).

Please share avalanche, snowpack or weather observations via our [website](#), email (mtavalanche@gmail.com), phone (406-587-6984), or Instagram (#gnfacobs).



Island Park

Near Island Park a person can trigger large to very large avalanches that break on buried weak layers. Dave was riding in the Centennials yesterday and found these weak layers on some slopes, but not all which can make assessment tricky. See Dave's [video](#) for information on what he found and what to do about it. Either avoid steep slopes or dig to look for and assess buried weak layer before entering steep terrain.

Upcoming Avalanche Education and Events

Our education calendar is full of awareness lectures and field courses. Check it out: [Events and Education Calendar](#).

January 4 + field day on January 7 or 8, *Avalanche Fundamentals for Snowmobilers*, Information and pre-registration [HERE](#).

January 4 + field day, *Avalanche Fundamentals for Skiers and Snowboarders*, Information and pre-registration [HERE](#).

Every Saturday, 10 a.m. - 2:00 p.m. *Avalanche Rescue Training*, drop in for any amount of time. Round Lake Warming Hut, Cooke City. Free.

[Loss in the Outdoors](#), is a support group for those who have been affected by grief and loss related to outdoor pursuits. Check out the link for more information.

Please consider donating to the Friends of GNFAC [Annual Fundraiser](#).